

Physica C 185–189 (1991) 1521–1522
North-Holland

PHYSICA C

CHARGE AND SPIN FLUCTUATIONS Cu^{2+} ($S=1/2$) \leftrightarrow Cu^{3+} ($S=1$) IN HTSCs. NEW MODEL OF NARROW IMPURITY BANDS

Mikhail EREMIN and Oleg ANIKEENOK

Physics Department, Kazan State University, 420008 Kazan, USSR

The energy band of oxide superconductors is theoretically investigated when charge and spin fluctuations of Cu^{2+} ($S=1/2$)- O^- ($S=0$) \leftrightarrow Cu^{3+} ($S=1$)- O^{2-} ($S=1/2$) take place. It is shown that two narrow bands arise near Fermi level under hole doping. The dynamic admixture of $3z^2-r^2$ state to the x^2-y^2 state is equal to approximately 10%.

The $\text{Cu}^{2+}3d^9$ configuration contains one hole in the 3d-shell. Extra holes in CuO_2 planes are added in HTSC materials and the equilibrium $\text{Cu}^{3+}\text{O}^{2-} \leftrightarrow \text{Cu}^{2+}\text{O}^-$ takes place¹. At present it is not clear what Cu^{3+} states are situated near Fermi level: high spin ($S=1$) or low spin ($S=0$) ones.

The goal of this paper is to calculate the structure of energy bands of normal phase CuO_2 planes assuming that high spin copper states are more favourable than low spin ones.

It is known that holes occupy copper states of x^2-y^2 symmetry in dielectric phase of metal oxides. Extra holes due to doping may occupy only states of the system with higher energy. These are either oxygen states or copper states². Let us write down the Hamiltonian as follows

$$H = \sum \varepsilon_1 a_{f\sigma}^+ a_{f\sigma} + \sum \varepsilon_2 b_{f\sigma}^+ b_{f\sigma} + \sum \varepsilon_j c_{j\sigma}^+ c_{j\sigma} \\ + \frac{1}{2} I_{11} \sum n_{f\sigma}^{(1)} n_{f\sigma}^{(1)} + I_{12} \sum n_{f\sigma}^{(1)} n_{f\sigma}^{(2)} \quad (1) \\ + K_{12} \sum a_{f\sigma}^+ b_{f\sigma}^+ a_{f\sigma} b_{f\sigma} + \sum t_{jl} c_{j\sigma}^+ c_{j'\sigma} \\ + \sum t_{fj}^{(1)} (a_{f\sigma}^+ c_{j\sigma} + c_{j\sigma}^+ a_{f\sigma}) + \sum t_{fj}^{(2)} (b_{f\sigma}^+ c_{j\sigma} + c_{j\sigma}^+ b_{f\sigma}).$$

Here 1 (2) - the energy of holes in copper states x^2-y^2 ($3z^2-r^2$) respectively, ε_j - the energy of oxygen holes with σ -symmetry, I_{11} (I_{22}) - the energy of Coulomb repulsion for holes in the states x^2-y^2 ($3z^2-r^2$), I_{12} and K_{12} - Coulomb and exchange integrals for holes in the states x^2-y^2 and $3z^2-r^2$, respectively.

Creation operators a^+ , b^+ and c^+ refer to holes in copper states x^2-y^2 , $3z^2-r^2$ and oxygen, respectively. Indices f and j numerate the sites of copper and oxygen.

It is convenient to analyse the energy bands starting from the case when $t^{(1)} = t^{(2)} = 0$. The energy operator for copper holes is diagonalised using Hubbard operators $X^{p,q}$. Eigenfunctions of the total hole spin are chosen as basis functions $|p\rangle$ and $|q\rangle$, for example

$$| \sigma_1, \sigma_2 \rangle = b_{\sigma_1}^+ a_{\sigma_2}^+ | v \rangle, \quad | 10 \rangle = \frac{1}{\sqrt{2}} \sum b_{\sigma_1}^+ a_{\sigma_2}^+ | v \rangle, \quad (2)$$

where $|v\rangle$ is the Cu^+ ground state.

There are eight Hubbard sub-bands. The energy sub-bands are given in the Table. The numerical estimates of energies are given relatively to the Hubbard sub-band E_3 . They are obtained based on the spectroscopic data³.

Table. Energy Hubbard sub-bands

$X^{p,q}$	Expression	E in eV
$X^{\sigma_2, \sigma_2 \bar{\sigma}_2}$	$E_8 = \varepsilon_2 + I_{22}$	4.3
$X^{\sigma_1, \sigma_1 \bar{\sigma}_1}$	$E_7 = \varepsilon_1 + I_{11}$	3.4
$X^{\sigma_1, 00}$	$E_6 = \varepsilon_2 + I_{12} + K_{12}$	3.4
$X^{\sigma_2, 00}$	$E_5 = \varepsilon_1 + I_{12} + K_{12}$	2.2
$X^{\sigma_1, \sigma_1, \sigma_2}, X^{\sigma_1, 10}$	$E_4 = \varepsilon_2 + I_{12} - K_{12}$	1
$X^{\sigma_2, \sigma_1 \sigma_2}, X^{\sigma_2, 10}$	$E_3 = \varepsilon_1 + I_{12} - K_{12}$	0
X^{v, σ_2}	$E_2 = \varepsilon_2$	-4.5
X^{v, σ_1}	$E_1 = \varepsilon_1$	-5.5